

THE LINDELÖF PROPERTY IN BANACH SPACES

STUDIA MATH. 154, 1-3, 165-192

B. CASCALES, I. NAMIOKA, AND J. ORIHUELA

ABSTRACT. A topological space (T, τ) is said to be fragmented by a metric d on T if each non-empty subset of T has non-empty relatively open subsets of arbitrarily small d -diameter. The basic theorem of the present paper is the following. Let (M, ρ) be a metric space with ρ bounded and let D be an arbitrary index set. Then for a compact subset K of the product space M^D the following four conditions are equivalent:

(i) K is fragmented by d_D , where, for each $S \subset D$,

$$d_S(x, y) = \sup\{\rho(x(t), y(t)) : t \in S\};$$

(ii) For each countable subset A of D , (K, d_A) is separable;

(iii) The space $(K, \gamma(D))$ is Lindelöf, where $\gamma(D)$ is the topology of uniform convergence on the family of countable subsets of D ;

(iv) $(K, \gamma(D))^{\mathbb{N}}$ is Lindelöf.

The rest of the paper is devoted to applications of the basic theorem. Here are some of them. A compact Hausdorff space K is Radon-Nikodým compact if, and only if, there is a bounded subset D of $C(K)$ separating the points of K such that $(K, \gamma(D))$ is Lindelöf. If X is a Banach space and H is a weak*-compact subset of the dual X^* which is weakly Lindelöf, then $(H, \text{weak})^{\mathbb{N}}$ is Lindelöf. Furthermore, under the same condition $\overline{\text{span}(H)}^{\|\cdot\|}$ and $\overline{\text{co}H}^{w^*}$ are weakly Lindelöf. The last conclusion answers a question by Talagrand. Finally we apply our basic theorem to certain classes of Banach spaces including weakly compactly generated ones and the duals of Asplund spaces.

1. INTRODUCTION

The starting point of the present investigation is a theorem by one of us in [25], namely, a Banach space X is an Asplund space if and only if its dual X^* is Lindelöf with respect to the topology of uniform convergence on bounded countable subsets of X , the γ -topology. In the present paper, we show that this result is a special case of a much more general theorem on function spaces and that it has interesting consequences including a solution to a question by Talagrand.

This paper is organized as follows. After the introduction of Section 1, the basic theorem and its important corollary are stated and proved in Section 2.

In Section 3, a new characterization of Radon-Nikodým compact spaces by the Lindelöf property relative to the γ -topology is derived from the basic theorem. It will be shown that Meyer's characterization of compact scattered spaces [23] by the Lindelöf property with respect to the G_δ -topology is also a consequence of the basic theorem.

1991 *Mathematics Subject Classification*. Primary 46B99, 46A50; Secondary 54C35.

Key words and phrases. Lindelöf spaces, Radon-Nikodým compact spaces, fragmentability, γ -topology.

Partially supported by the research grant DGES PB 98-0381.

In Section 4, we use the Lindelöf property relative to the γ -topology to study weakly Lindelöf property of sets in dual Banach spaces. We show, for instance, that the weak*-closed convex hull of a weak*-compact subset which is weakly Lindelöf in a dual Banach space is again weakly Lindelöf. This solves a problem of Talagrand in [32].

The theme of Section 4 is further expanded in Section 6 where it is proved, in particular, that the norm closed linear span of a weak*-compact subset in a dual Banach space that is weakly Lindelöf is a WLD Banach space, as defined in Section 6. It should be noted here that each WLD Banach space is weakly Lindelöf and more. Our approach depends on the existence of “projectional generators” shown in Section 5. Our results on projectional generators also give a unified approach to the existence of projectional resolutions of the identity for both weakly compactly generated Banach spaces and dual of Asplund spaces.

In Section 7, we present several examples that illustrate the results of Sections 5 and 6

Our notation and terminology are standard and we take the books by Engelking and Kelley, [10] and [20], as our references for topology. Compact spaces are usually referred to by letters K, H, \dots and our normed spaces X, Y, \dots are assumed to be real. Given a topological space Z we let $C(Z)$ (resp. $C_b(Z)$) denote the space of real continuous (resp. real continuous uniformly bounded) functions defined on Z . Given a Banach space X , B_X denotes its closed unit ball and X^* denotes its dual space. When F is a subset of X^* , we write $\sigma(X, F)$ to denote the locally convex topology (maybe non-Hausdorff) on X of pointwise convergence on F ; $\sigma(X, X^*)$ is the weak topology of X and $\sigma(X^*, X)$ is the weak* topology of X^* . $C_b(Z)$ will be considered as Banach space endowed with the supremum norm.

2. FRAGMENTABILITY AND THE LINDELÖF PROPERTY FOR $\gamma(D)$

We first gather definitions of the terms and notation necessary for stating the main theorem of the present paper, Theorem 2.1. Recall that a topological space is said to be *Lindelöf* if each open cover of the space admits a countable subcover. The following definition is due to Jayne and Rogers [19].

Definition 1. *Let (Z, τ) be a topological space and ρ a metric on Z . We say that (Z, τ) is fragmented by ρ (or ρ -fragmented) if for each non-empty subset C of Z and for each $\varepsilon > 0$ there exists a non-empty τ -open subset U of Z such that $U \cap C \neq \emptyset$ and $\rho\text{-diam}(U \cap C) \leq \varepsilon$.*

It is easily checked that for (Z, τ) to be ρ -fragmented, it is sufficient that each τ -closed non-empty subset of X has non-empty relatively τ -open subsets of arbitrarily small ρ -diameter.

Let (M, ρ) be a metric space and let D be an arbitrary set. We shall write $\tau_p(D)$ (only τ_p if no ambiguity is likely) to denote the product topology of the space M^D . Assume henceforth that ρ is bounded, which can always be done without altering the uniformity of M . For any set $S \subset D$ we define the pseudo-metric d_S on M^D by the formula

$$(1) \quad d_S(x, y) = \sup\{\rho(x(t), y(t)) : t \in S\}, \quad \text{for } x, y \in M^D.$$

The metric d_D will be simply denoted by d ; the topology associated to d in M^D is the topology of uniform convergence on D . Let $\gamma(D)$ denote the uniform topology

on M^D generated by the family of pseudo-metrics $\{d_A : A \subset D, A \text{ countable}\}$, i.e. the topology of uniform convergence on the family of countable subsets of D .

The following notation is used in the proof of the next theorem. Let $2^{\mathbb{N}}$ be the space of all sequences of 0's and 1's and let $2^{(\mathbb{N})}$ be the set of all finite sequences of 0's and 1's. For a given $t \in 2^{(\mathbb{N})}$, let $|t|$ denote the length of t ; for $\sigma \in 2^{\mathbb{N}}$ and $n \in \mathbb{N}$, we write $\sigma|n = (\sigma(1), \dots, \sigma(n)) \in 2^{(\mathbb{N})}$.

Theorem 2.1. *Let (M, ρ) and D be as above, and let K be a compact subset of (M^D, τ_p) . Then the following conditions are equivalent:*

- (a) *The space (K, τ_p) is fragmented by d ;*
- (b) *For each countable subset A of D , (K, d_A) is separable;*
- (c) *The space $(K, \gamma(D))$ is Lindelöf.*

Proof.- (a) \Rightarrow (b) By Lemma 2.1 of [24], $(K|_A, \tau_p(A))$ is fragmented by d_A . Since M^A is metrizable, $(K|_A, \tau_p(A))$ is compact metrizable; hence it has a countable base. If (K, d_A) is not separable, then there is an uncountable subset Q of $K|_A$ and $\varepsilon > 0$ such that $d_A(p, q) > \varepsilon$ whenever $p, q \in Q$ and $p \neq q$. We may assume that no point of Q is τ_p -isolated in Q since $(K|_A, \tau_p(A))$ has a countable base. Since $(K|_A, \tau_p(A))$ is fragmented by d_A , there is a $\tau_p(A)$ -open subset U of $K|_A$ such that $U \cap Q \neq \emptyset$ and d_A -diam $(U \cap Q) < \varepsilon$. Hence $U \cap Q$ is a singleton, contradicting the fact that no point of Q is $\tau_p(A)$ -isolated in Q . Hence (K, d_A) is separable.

(b) \Rightarrow (a) Suppose that (K, τ_p) is not fragmented by d . Then, for some non-empty τ_p -closed subset C of K and $\varepsilon > 0$, each non-empty τ_p -open subset of C has d -diameter greater than ε . By induction on $n = |s|$, $s \in 2^{(\mathbb{N})}$, we construct a family $\{U_s : s \in 2^{(\mathbb{N})}\}$ of non-empty relatively τ_p -open subsets of C and a family $\{t_s : s \in 2^{(\mathbb{N})}\}$ of points of D , satisfying the following conditions:

- (α) $U_\emptyset = C$,
- (β) for each s , $\overline{U_{s0}}^{\tau_p} \cup \overline{U_{s1}}^{\tau_p} \subset U_s$,
- (γ) $\rho(x(t_s), y(t_s)) > \varepsilon$ for each $x \in \overline{U_{s0}}^{\tau_p}$ and $y \in \overline{U_{s1}}^{\tau_p}$.

Construction. (α) starts the induction from $n = 0$. Next, for some $n > 0$, assume that $\{U_s : |s| < n\}$ and $\{t_s : |s| < n - 1\}$ have been constructed. Fix an $s \in 2^{(\mathbb{N})}$ with $|s| = n - 1$. By hypothesis, there are $x, y \in U_s$, with $d(x, y) > \varepsilon$. Hence for some $t_s \in D$, $\rho(x(t_s), y(t_s)) > \varepsilon$. By the τ_p -continuity of the map

$$(x', y') \mapsto \rho(x'(t_s), y'(t_s))$$

there are relatively τ_p -open neighborhoods U_{s0} and U_{s1} of x and y , respectively, so that (β) and (γ) are satisfied. This completes the construction. Note that (γ) implies that $\overline{U_{s0}}^{\tau_p} \cap \overline{U_{s1}}^{\tau_p} = \emptyset$ for each $s \in 2^{(\mathbb{N})}$.

For each $\sigma \in 2^{\mathbb{N}}$, choose $x_\sigma \in \bigcap_{n=1}^{\infty} \overline{U_{\sigma|n}}^{\tau_p}$. If $\sigma, \sigma' \in 2^{\mathbb{N}}$ are two different sequences, then for some $n \in \{0\} \cup \mathbb{N}$, $\sigma|n = \sigma'|n$ and $\sigma|(n + 1) \neq \sigma'|(n + 1)$. Then by (γ) we have $\rho(x_\sigma(t_{\sigma|n}), x_{\sigma'}(t_{\sigma|n})) > \varepsilon$. Letting $A = \{t_s : s \in 2^{(\mathbb{N})}\}$ we have $d_A(x_\sigma, x_{\sigma'}) > \varepsilon$. Since $2^{\mathbb{N}}$ is uncountable, (K, d_A) is not separable, and therefore (b) does not hold.

(c) \Rightarrow (b) This is clear because the topology associated to d_A is weaker than $\gamma(D)$ whenever A is a countable subset of D .

(a) & (b) \Rightarrow (c) Let $\mathcal{U} = \{U_j : j \in J\}$ be a $\gamma(D)$ -open cover of K and let $\mathcal{C} = \{A : A \subset D \text{ and } A \text{ is countable}\}$. Without loss of generality we may assume that each U_j is of the form

$$U_j = U(x_j, A_j, \varepsilon_j) \stackrel{\text{def}}{=} \{y \in K : d_{A_j}(x_j, y) < \varepsilon_j\},$$

where $x_j \in K$, $A_j \in \mathcal{C}$ and $\varepsilon_j > 0$. For each $A \in \mathcal{C}$, let us define

$$\mathcal{U}(A) = \{U_j : j \in J, A_j \subset A\} \text{ and } U(A) = \bigcup \{U_j : U_j \in \mathcal{U}(A)\}.$$

Then we have

$$(2) \quad \mathcal{U} = \bigcup \{\mathcal{U}(A) : A \in \mathcal{C}\} \text{ and } K = \bigcup \{U(A) : A \in \mathcal{C}\}.$$

Also if $A \subset A'$ then $U(A) \subset U(A')$.

We **claim** that $K = U(A)$ for some $A \in \mathcal{C}$.

Suppose for a moment this is true. Then since each member of $\mathcal{U}(A)$ is d_A -open and since (K, d_A) is separable by (b), there is a countable subfamily of $\mathcal{U}(A)$ (hence of \mathcal{U}) that covers K , which completes the proof.

The proof of the claim is by contradiction. So assume that $U(A) \neq K$ for each $A \in \mathcal{C}$. For each $A \in \mathcal{C}$, let

$$C(A) = K \setminus U(A) \text{ and } C = \bigcap \{\overline{C(A)}^{\tau_p} : A \in \mathcal{C}\}.$$

We note that $C(A) \supset C(A')$ whenever $A \subset A'$. By compactness of (K, τ_p) , $C \neq \emptyset$, and now (a) tells us that (C, τ_p) is fragmented by d . So by Lemma 1.1 of [24], there is a point $y \in C$ where the identity map $(C, \tau_p) \rightarrow (C, d)$ is continuous. The second equality in (2) ensures us that $y \in U(B)$ for some $B \in \mathcal{C}$. Since $U(B)$ is d_B -open, for some $\varepsilon > 0$, $y \in U(y, B, \varepsilon) \subset U(B)$. Then for each $x \in C(B) = K \setminus U(B)$, $x \notin U(y, B, \varepsilon)$ and so for some $t \in B$, $\rho(x(t), y(t)) \geq 2\varepsilon/3$. For each $t \in B$, let

$$(3) \quad D_t = \{x \in C(B) : \rho(x(t), y(t)) \geq 2\varepsilon/3\}.$$

Then from above $C(B) = \bigcup \{D_t : t \in B\}$.

Let V be a τ_p -open neighborhood of y in K such that $d\text{-diam}(\overline{V}^{\tau_p} \cap C) \leq \varepsilon/2$. Then we claim that, for some $t \in B$, $D_t \cap V \cap C(A) \neq \emptyset$ for each $A \in \mathcal{C}$. For, otherwise, for each $t \in B$ there is an $A_t \in \mathcal{C}$ such that $D_t \cap V \cap C(A_t) = \emptyset$. Since B is countable, the set $E \stackrel{\text{def}}{=} B \cup \bigcup \{A_t : t \in B\}$ is also countable, and $D_t \cap V \cap C(E) = \emptyset$ for all $t \in B$. Hence

$$\emptyset = \left(\bigcup \{D_t : t \in B\} \right) \cap V \cap C(E) = C(B) \cap V \cap C(E) = V \cap C(E),$$

contradicting $y \in C \subset \overline{C(E)}^{\tau_p}$.

Now fix a $t \in B$ so that $D_t \cap V \cap C(A) \neq \emptyset$ for each $A \in \mathcal{C}$, and let

$$z \in \bigcap \{\overline{D_t \cap V \cap C(A)}^{\tau_p} : A \in \mathcal{C}\}.$$

Then $z \in \overline{V}^{\tau_p} \cap C$, and so

$$(4) \quad d(z, y) \leq \varepsilon/2.$$

On the other hand, since $z \in \overline{D_t}^{\tau_p}$, it follows by (3) that $\rho(z(t), y(t)) \geq 2\varepsilon/3$, which contradicts (4). This completes the proof both of the claim and of the theorem. ■

It is well known that the product of two Lindelöf spaces is not in general Lindelöf again: indeed let $Z = \mathbb{R}$ and endow it with the topology for which a basis is given by all the intervals $[x, r)$, where $x, r \in \mathbb{R}$, $x < r$ and r is a rational number; then Z is a separable first-countable space that is Lindelöf and which is not second-countable; moreover $Z \times Z$ is not normal and therefore not Lindelöf, see [10, pages 248-249].

Fortunately the Lindelöf property for the spaces $(K, \gamma(D))$ in Theorem 2.1 is preserved under the countable power.

Corollary 2.2. *Let K, M, D be as in Theorem 2.1. If K satisfies one of the three conditions of the theorem, then $(K, \gamma(D))^{\mathbb{N}}$ is Lindelöf. In particular, $(K, \gamma(D))^n$ is Lindelöf for each $n \in \mathbb{N}$.*

Proof.- We may assume that the metric ρ of the space M is bounded by 1. Let $\varphi : (M^D)^{\mathbb{N}} \rightarrow (M^{\mathbb{N}})^D$ be the map defined by $\varphi(\xi)(t)(j) = \xi(j)(t)$ for all $\xi \in (M^D)^{\mathbb{N}}, t \in D, j \in \mathbb{N}$. Clearly φ is a homeomorphism when the product topology is used throughout. Now the space $M^{\mathbb{N}}$ is metrizable, and we use the metric $\rho_{\infty}(m, m') \stackrel{\text{def}}{=} \sum_{j \in \mathbb{N}} 2^{-j} \rho(m(j), m'(j))$ for $m, m' \in M^{\mathbb{N}}$. Let d_{∞} be the metric on $(M^{\mathbb{N}})^D$ given by

$$d_{\infty}(x, x') \stackrel{\text{def}}{=} \sup\{\rho_{\infty}(x(t), x'(t)) : t \in D\} \text{ for } x, x' \in (M^{\mathbb{N}})^D.$$

We now show that if K is fragmented by d then $\varphi(K^{\mathbb{N}})$ is fragmented by d_{∞} . Let $\varepsilon > 0$, let C be a non-empty subset of $K^{\mathbb{N}}$ and let $\pi_i : K^{\mathbb{N}} \rightarrow K$ be the i -th projection. Then by induction we can construct a decreasing sequence $V_1 \supset V_2 \supset \dots$ of non-empty relatively open subsets of C such that $d\text{-diam } \pi_j(V_j) < \varepsilon/2$ for each $j \in \mathbb{N}$. Choose $k \in \mathbb{N}$ so that $2^{-k} < \varepsilon/2$, and let $\xi, \xi' \in V_k$. Then for each $t \in D$,

$$\begin{aligned} \rho_{\infty}(\varphi(\xi)(t), \varphi(\xi')(t)) &\leq \sum_{j \leq k} 2^{-j} \rho(\xi(j)(t), \xi'(j)(t)) + \sum_{j \geq k+1} 2^{-j} \\ &< \sum_{j \leq k} 2^{-j} d(\pi_j(\xi), \pi_j(\xi')) + \varepsilon/2 \leq \varepsilon/2 + \varepsilon/2 = \varepsilon. \end{aligned}$$

Thus $\varphi(V_k)$ is a non-empty relatively open subset of $\varphi(C)$ with ρ_{∞} -diameter not greater than ε .

Hence by Theorem 2.1, $\varphi(K^{\mathbb{N}})$ is $\gamma(D)$ -Lindelöf. So we finish the proof by showing that φ maps $(M^D, \gamma(D))^{\mathbb{N}}$ homeomorphically onto $((M^{\mathbb{N}})^D, \gamma(D))$. Let τ_1, τ_2 be the topologies of these two spaces respectively. Then a net ξ_{α} in $(M^D)^{\mathbb{N}}$ τ_1 -converges to $\xi \in (M^D)^{\mathbb{N}}$ if and only if, (i) for each $j \in \mathbb{N}$ and for each countable set $A \subset D$, $\rho(\xi_{\alpha}(j)(t), \xi(j)(t)) \rightarrow 0$ uniformly in $t \in A$. On the other hand, the net $\varphi(\xi_{\alpha})$ τ_2 -converges to $\varphi(\xi)$ if and only if, (ii) for each countable $A \subset D$,

$$\rho_{\infty}(\varphi(\xi_{\alpha})(t), \varphi(\xi)(t)) = \sum_{j \in \mathbb{N}} 2^{-j} \rho(\xi_{\alpha}(j)(t), \xi(j)(t)) \rightarrow 0$$

uniformly in $t \in A$. The equivalence of statements (i) and (ii) can be seen by an easy calculation similar to the one given above. Hence φ is a τ_1 - τ_2 homeomorphism. ■

As an immediate consequence of the foregoing, we obtain the following theorem, the first part of which was mentioned in the introduction. It has been stated in

[25] as Theorem B and C. The original proof is quite different and depends on the technique of projections in Banach spaces.

Theorem 2.3. ([25]) *A Banach space X is an Asplund space if and only if $(X^*, \gamma(B_X))$ is Lindelöf. If this is the case, then $(X^*, \gamma(B_X))^n$ is Lindelöf for each $n \in \mathbb{N}$.*

Proof.- Note that $(X^*, \gamma(B_X))^n$ is Lindelöf if and only if $(B_{X^*}, \gamma(B_X))^n$ is Lindelöf, and X is an Asplund space if and only if (B_{X^*}, weak^*) is fragmented by the norm. Therefore the theorem follows from 2.1 and 2.2 by regarding (B_{X^*}, weak^*) as a compact subspace of $([-1, 1]^{B_X}, \tau_p)$. ■

3. APPLICATIONS TO RN-COMPACT SPACES

Let K be a compact Hausdorff space and let D be a uniformly bounded subset of $C(K)$ and $A \subset D$, then we define the pseudo-metric on K by

$$d_A(x, x') = \sup\{|f(x) - f(x')| : f \in A\}, \quad \text{for } x, x' \in K.$$

We write again $\gamma(D)$ to denote the uniform topology on K generated by the family of pseudo-metrics $\{d_A : A \subset D, A \text{ countable}\}$. Observe that when D separates the points of K , K embeds in $[-m, m]^D$ for some $m > 0$. Hence the topology $\gamma(D)$ now defined is the one already given through the embedding $K \subset [-m, m]^D$, and $\gamma(D)$ is stronger than the original topology of K . In particular the equivalences we have seen in Theorem 2.1 and Corollary 2.2 remain true.

Theorem 3.1. *Let K be a compact Hausdorff space and let D be a uniformly bounded subset of $C(K)$. Then the following statements are equivalent.*

- (i) *The space (K, d_A) is separable for each countable $A \subset D$;*
- (ii) *The space $(K, \gamma(D))$ is Lindelöf;*
- (iii) *The space $(K, \gamma(D))^{\mathbb{N}}$ is Lindelöf.*

Proof.- From the remark above, the theorem is clear in case D separates the points of K . The general case can be reduced to this as follows. Let $m = \sup\{\|f\| : f \in D\}$ and let $\varphi : K \rightarrow [-m, m]^D$ be the map given by $\varphi(x)(f) = f(x)$ for all $x \in K$ and $f \in D$. Then $K' \stackrel{\text{def}}{=} \varphi(K)$ is a compact Hausdorff space. For each $f \in D$, let $\hat{f} \in C(K')$ be the map given by $\hat{f}(\varphi(x)) = f(x)$, and, for each $A \subset D$, let $\hat{A} = \{\hat{f} : f \in A\}$. Then clearly $f \mapsto \hat{f}$ is a one-to-one map of D onto \hat{D} and $d_A(x, y) = d_{\hat{A}}(\varphi(x), \varphi(y))$ for all $x, y \in K$. It follows that (K, d_A) is separable if, and only if, $(K', d_{\hat{A}})$ is separable. The last equality also implies that, for each $x \in K$, $\{y \in K : d_A(x, y) < \varepsilon\} = \varphi^{-1}(\{z \in K' : d_{\hat{A}}(\varphi(x), z) < \varepsilon\})$. Hence a subset U of K is $\gamma(D)$ -open if and only if $U = \varphi^{-1}(U')$ for some $\gamma(\hat{D})$ -open subset U' of K' . From this it is straight forward to check that $(K, \gamma(D))$ (resp. $(K, \gamma(D))^{\mathbb{N}}$) is Lindelöf if, and only if, $(K', \gamma(\hat{D}))$ (resp. $(K', \gamma(\hat{D}))^{\mathbb{N}}$) is Lindelöf. Since \hat{D} separates the points of K' , the conclusion of the theorem is true for \hat{D} and K' . Hence the theorem is proved in general. ■

A compact Hausdorff space is said to be *Radon-Nikodým compact* (or RN compact) if it is homeomorphic to a weak*-compact subset of the dual of an Asplund space, *i.e.* a dual Banach space with the RNP. It is shown in [24] that a compact Hausdorff space is RN compact if and only if it is fragmented by a lower-semicontinuous metric on the space. When (M, ρ) is a metric space (with ρ

bounded) the metric d in Theorem 2.1 is clearly τ_p lower semi-continuous. Therefore, Theorem 2.1 provides the following characterization of RN-compact spaces.

Proposition 3.2. *A compact Hausdorff space is RN compact if, and only if, it is homeomorphic to a pointwise compact subset K of $[-1, 1]^D$, for some set D , such that $(K, \gamma(D))$ is Lindelöf.*

Proof.- By Theorem 3.6 of [24] a compact space is RN compact if, and only if, K is homeomorphic to a pointwise compact subset K of $[-1, 1]^D$, for some set D , such that (K, d_A) is separable for each countable subset A of D . An application of Theorem 2.1 finishes the proof of the proposition. ■

In terms of spaces of continuous functions the proposition above can be restated as follows.

Corollary 3.3. *A compact Hausdorff space K is RN compact if, and only if, there is a bounded subset D of $C(K)$ separating points of K such that $(K, \gamma(D))$ is Lindelöf. Moreover, if this is the case, then $(K, \gamma(D))^{\mathbb{N}}$ is Lindelöf.*

Proof.- Assume K is RN compact. By Proposition 3.2, we may assume that K is a subspace of $([-1, 1]^D, \tau_p)$ for certain set D , with $(K, \gamma(D))$ Lindelöf; for every $d \in D$ let $\pi_d : [-1, 1]^D \rightarrow [-1, 1]$ be the projection defined by $\pi_d(x) = x(d)$, $x \in [-1, 1]^D$. If we let as $D = \{\pi_d : d \in D\}$, then D is a uniformly bounded subset of $C(K)$ separating the points of K and such that $(K, \gamma(D))$ is Lindelöf. The last part follows from Theorem 3.1. Similar argument proves the converse. ■

For weakly compact subsets of $C(K)$, we have the following.

Corollary 3.4. *Let K be a compact Hausdorff space and let $H \subset C(K)$ be a weakly compact (i.e. bounded and τ_p -compact) set. Then $(K, \gamma(H))^{\mathbb{N}}$ is Lindelöf.*

Proof.- Given a countable $A \subset H$, $\overline{A}^{\tau_p} \subset C(K)$ is $\tau_p(K)$ -metrizable and thus the space $(C(\overline{A}^{\tau_p}), d_{\overline{A}^{\tau_p}})$ is separable. Hence, $(K|_{\overline{A}^{\tau_p}}, d_{\overline{A}^{\tau_p}})$ is separable and (K, d_A) too. In view of Theorem 3.1, the proof is complete. ■

We need the following easy lemma that appears in [5] in a more general context.

Lemma 1. *Let Z be a Lindelöf space, and let $H \subset C(Z)$ be equicontinuous. Then $(H, \tau_p(Z))$ is metrizable.*

Proof. Let d_H be the pseudo-metric on Z given by

$$d_H(z, z') = \min\{1, \sup_{h \in H} |h(z) - h(z')|\}.$$

Since H is equicontinuous, the d_H -topology is weaker than the given one on Z . So (Z, d_H) is Lindelöf and hence separable. Let D be a countable d_H -dense subset of Z . Then since H is d_H -equicontinuous, on H the topologies of pointwise convergence on D and on Z coincide. Therefore $(H, \tau_p(Z))$ is metrizable. ■

Given a subset D of \mathbb{R}^K , let

$$F(D) = \bigcup \{\overline{A}^{\tau_p} : A \subset D, A \text{ countable}\}.$$

Note that if B is a countable subset of $F(D)$ then there is a countable subset A of D such that $\overline{B}^{\tau_p} \subset \overline{A}^{\tau_p} \subset F(D)$. In particular, $F(F(D)) = F(D)$.

Recall that a topological space Z is said to be *countably tight* (resp. to be a *Fréchet-Urysohn space*) if for each set $S \subset Z$ and each point $x \in \overline{S}$ there is

countable set $A \subset S$ (resp. sequence $(x_n)_n$ in S) such that $x \in \bar{A}$ (resp. $(x_n)_n$ converges to x), see [2, pages 5 and 7]. In applying the results of the last section, the following theorem of Arkhangel'skii ([2, Theorem II.1.1]) is very useful. We quote a special case.

Theorem A. *Let T be a topological space such that T^n is Lindelöf for each $n \in \mathbb{N}$. Then $(C(T), \tau_p(T))$ is countably tight.*

Corollary 3.5. *Let K be a compact space and let D be a bounded subset of $C(K)$ such that $(K, \gamma(D))$ is Lindelöf. Then the following properties hold:*

- (a) *For any countable set $A \subset D$, \bar{A}^{τ_p} (closure taken in \mathbb{R}^K) is $\gamma(D)$ -equicontinuous and τ_p -metrizable;*
- (b) *$F(D) = C(K, \gamma(D)) \cap \bar{D}^{\tau_p}$, where the closure is taken in \mathbb{R}^K ;*
- (c) *$(F(D), \tau_p)$ is a Fréchet-Urysohn space.*

Proof.- (a) easily follows from the previous lemma: if $A \subset D$ is countable then A is $\gamma(D)$ -equicontinuous; its τ_p -closure \bar{A}^{τ_p} in \mathbb{R}^K is again $\gamma(D)$ -equicontinuous and therefore τ_p -metrizable after Lemma 1. This proves (a)

For (b), we first note that (a) implies $F(D) \subset C(K, \gamma(D)) \cap \bar{D}^{\tau_p}$. Next we note that $(K, \gamma(D))^n$ is Lindelöf for each $n \in \mathbb{N}$ by Theorem 3.1. This fact implies that $(C(K, \gamma(D)), \tau_p)$ is countably tight according to Theorem A. Therefore if $f \in C(K, \gamma(D)) \cap \bar{D}^{\tau_p}$ then there is a countable subset A of D such that $f \in \bar{A}^{\tau_p}$. Hence $f \in F(D)$ which proves (b). The proof of (c) is similar: Suppose that $S \subset F(D)$ and $f \in \bar{S}^{\tau_p} \cap F(D)$. Then by the countable tightness, there is a countable subset B of S such that $f \in \bar{B}^{\tau_p}$. Then as noted above, there is a countable subset A of D such that $\bar{B}^{\tau_p} \subset \bar{A}^{\tau_p}$. In particular \bar{B}^{τ_p} is τ_p -metrizable by (a). Therefore there is a sequence in B (hence in S) that τ_p -converges to f . This proves (c). ■

Recall that a topological space T is said to be *scattered* if each non-empty subset of T has an isolated point, or equivalently T is fragmented by the (necessarily lower-semicontinuous) trivial metric ρ , where $\rho(t, s) = 0$ for $t = s$ and $\rho(t, s) = 1$ for $t \neq s$. It can be shown (cf. [30, Theorem 8.5.4]) that a compact Hausdorff space K is scattered if and only if there is no continuous map from K onto $[0, 1]$. We remark that in the corollary above if $B_{C(K)} \subset F(D)$ then K is scattered. For then, $(B_{C(K)}, \tau_p)$ is a Fréchet-Urysohn space; on the other hand, $(B_{C[0,1]}, \tau_p)$ is not Fréchet-Urysohn, see [2, Lemma II.3.5], and consequently K cannot be continuously mapped onto $[0, 1]$.

Given a topological space (Z, \mathcal{T}) , the G_δ -topology associated to \mathcal{T} is the topology on Z whose basis is the family of G_δ -sets, $\{\bigcap_n U_n : U_n \in \mathcal{T}\}$; when no confusion is likely we simply write Z for the topological space and then refer to its G_δ -topology.

Lemma 2. *Let K be a compact Hausdorff space. Then the G_δ -topology for K is identical with $\gamma(B_{C(K)})$ on K .*

Proof.- Clearly the G_δ -topology is stronger than $\gamma(B_{C(K)})$. Let $a \in K$, and let G be a G_δ -set containing a . Then $G = \bigcap_{n=1}^\infty U_n$ where each U_n is open in K . For each n , let f_n be a continuous function $f_n : K \rightarrow [0, 1]$ such that $f_n(a) = 0$, and $f_n|_{K \setminus U_n} \equiv 1$. Write $A = \{f_n : n \in \mathbb{N}\}$. Then A is a countable subset of $B_{C(K)}$,

and $x \in G$ whenever $d_A(a, x) < 1$, i.e.

$$a \in \{x \in K : d_A(a, x) < 1\} \subset G.$$

This shows that $\gamma(B_{C(K)})$ is stronger than the G_δ -topology and we are done. ■

Corollary 3.6 (Meyer, [23]). *For a compact Hausdorff space K , let τ_δ denote its G_δ -topology. Then the following statements are equivalent:*

- (a) K is scattered;
- (b) (K, τ_δ) is Lindelöf.
- (c) $(B_{C_b(K, \tau_\delta)}, \tau_p)$ is a Fréchet-Urysohn space.

Proof.- (a) \Leftrightarrow (b) Regarding K as a subset of $([-1, 1]^{B_{C(K)}}, \tau_p)$, we apply Theorem 2.1. In this case the metric d is twice the trivial metric and the topology $\gamma(B_{C(K)})$ is the G_δ -topology for K by the lemma above. (a) \Leftrightarrow (b) now follows.

Next assume (b), and we apply Corollary 3.5 to our K and $D \stackrel{\text{def}}{=} B_{C(K)}$. The hypotheses are satisfied by (b). Since the τ_p -closure of D is $[-1, 1]^K$, (b) of Corollary 3.5 says that $F(D) = B_{C_b(K, \gamma(D))} = B_{C_b(K, \tau_\delta)}$ and (c) of the same corollary says that $(B_{C_b(K, \tau_\delta)}, \tau_p)$ is a Fréchet-Urysohn space. This is (c). If (c) holds, then $(B_{C(K)}, \tau_p)$ is also a Fréchet-Urysohn space. But as remarked above, this implies (a). ■

We should comment here that topological spaces for which G_δ -sets are again open are called P -spaces. It is a very easy exercise to prove that if Z is a Lindelöf P -space then Z^n is Lindelöf for $n \in \mathbb{N}$ and so $(C(Z), \tau_p)$ has countable tightness; it also follows from Lemma 1 that for such a Z the separable subsets of $(C(Z), \tau_p)$ are metrizable, and hence $(C(Z), \tau_p)$ is Fréchet-Urysohn, see also [2]. Our argument also shows the fact that, for K compact and scattered, the space of all continuous functions on K endowed with its G_δ -topology is $B_1(K)$, the space of τ_p -limits of sequences in $C(K)$, and that all classes of Baire functions on K are the same [22].

4. POINTWISE LINDELÖF SUBSETS OF SPACES OF CONTINUOUS FUNCTIONS

Let D be a dense subset of a compact Hausdorff K and let H be bounded and $\tau_p(D)$ -compact subset of $C(K)$. In this section, we investigate the $\tau_p(K)$ -Lindelöf property of H by means of the $\gamma(D)$ -topology of the earlier sections. As application we can prove the results mentioned in the introduction.

The following simple proposition enables us to extract information on $(H, \tau_p(K))$ from that on $(H, \gamma(D))$.

Proposition 4.1. *Let K be a compact Hausdorff space, D a dense subset K and H a subset $C(K)$. If H is $\tau_p(K)$ -Lindelöf, then $\gamma(D)$ is stronger than $\tau_p(K)$ on H .*

Proof.- Let $f \in H$, $\varepsilon > 0$, $x \in K$, and

$$U = \{g \in H : |g(x) - f(x)| < \varepsilon\}.$$

Then U is a $\tau_p(K)$ -open neighborhood of f in H , and it is sufficient to show that U is a $\gamma(D)$ -neighborhood of f in H . For each $d \in D$, let

$$D_d = \{g \in H : |g(d) - f(d)| \leq \varepsilon/2\}.$$

If $g \in \bigcap \{D_d : d \in D\}$, then $|g(x) - f(x)| \leq \varepsilon/2$ since $x \in \overline{D}$, and therefore $g \in U$. It follows that $\bigcap \{D_d : d \in D\} \subset U$. Since each D_d is $\tau_p(K)$ -closed and H is $\tau_p(K)$ -Lindelöf, there is a countable subset A of D such that already $\bigcap \{D_d : d \in A\} \subset U$, i.e. $\{g \in H : \sup_{d \in A} |g(d) - f(d)| \leq \varepsilon/2\} \subset U$. Hence U is a $\gamma(D)$ -neighborhood of f in H and the proof is finished. ■

Corollary 4.2. *Let K be a compact Hausdorff space, D a dense subset K and H a bounded $\tau_p(D)$ -compact subset of $C(K)$. If $(H, \tau_p(K))$ is Lindelöf, then $(H, \tau_p(K))^{\mathbb{N}}$ is Lindelöf.*

Proof.- If H is $\tau_p(D)$ -compact and $\tau_p(K)$ -Lindelöf, then by [4, Theorem B] H is fragmented by the supremum norm of $C(K)$, i.e. as a compact subset H of $[-m, m]^D$ for a suitable m , H is fragmented by d in the notation of Theorem 2.1. According to Theorem 2.1 and Corollary 2.2, $(H, \gamma(D))^{\mathbb{N}}$ is Lindelöf. By Proposition 4.1, $\gamma(D)$ is stronger than $\tau_p(K)$ on H and therefore $(H, \tau_p(K))^{\mathbb{N}}$ is Lindelöf because it is a continuous image of the Lindelöf space $(H, \gamma(D))^{\mathbb{N}}$. ■

In [2, Problem IV.11.11] Arkhangel'skii asks the following question. Let K be a compact Hausdorff space. If there exists a τ_p -Lindelöf subset H of $C(K)$ that separates the points of K , is K countably tight? The next corollary is an answer to this question under a rather strong restriction on H .

Corollary 4.3. *Let K be a compact Hausdorff space, and H a $\tau_p(K)$ -Lindelöf bounded subset of $C(K)$ separating the points of K . If H is $\tau_p(D)$ -compact for some dense subset $D \subset K$, then K is countably tight.*

Proof.- An application of Corollary 4.2 allows us to conclude that $(H, \tau_p(K))^n$ is Lindelöf for $n \in \mathbb{N}$. Hence the space $C(H, \tau_p(K))$ is countably tight by Theorem A. The space K is homeomorphic to a subspace of $C(H, \tau_p(K))$ because H separates the points of K , and so the proof is done. ■

If X is a Banach space, then $B_{X^{**}}$ is always assumed to have the weak*-topology ($= \sigma(X^{**}, X^*)$) unless other topology is specified. Also X and B_X are considered as subspace/subset of X^{**} and $B_{X^{**}}$, respectively, by means of the canonical embedding. Thus (X^*, weak^*) is a subspace of $(C(B_{X^{**}}), \tau_p(B_X))$ and (X^*, weak) is a subspace of $(C(B_{X^{**}}), \tau_p(B_{X^{**}}))$. For a subset S of X^* , the weak and weak* closures of S are respectively denoted by \overline{S}^w and \overline{S}^{w^*} . A particular case of Corollary 4.2 is the following:

Corollary 4.4. *Let X be a Banach space and let H be a weak*-compact subset of X^* which is weakly Lindelöf. Then, $(H, \text{weak})^{\mathbb{N}}$ is Lindelöf.*

The next result gives the positive answer to a question posed by Talagrand that appears in [32] as *Problème 4.5*.

Theorem 4.5. *Let X be a Banach space and let H be a weak*-compact subset of X^* which is weakly Lindelöf. Then,*

- (a) $\overline{\text{co}(H)}^{w^*} = \overline{\text{co}(H)}^{\|\cdot\|}$;
- (b) $\overline{\text{co}(H)}^{w^*}$ is weakly Lindelöf.

Proof.- If H is a weak*-compact subset of X^* which is also weakly Lindelöf, then (H, weak^*) is fragmented by the dual norm by Corollary E in [4]. The equality in item (a) follows now from Theorem 2.3 in [24].

Let us prove (b). As noted in the proof of (a), (H, weak^*) is fragmented by the norm. Therefore if we let $W = \overline{\text{co}(H)}^{w^*}$, then W is weak*-compact and (W, weak^*) is fragmented by the norm by [24, Theorem 2.5]. By embedding W into $[-m, m]^{B_X}$ for a suitable $m > 0$, we see that $(W, \gamma(B_X))$ is Lindelöf by Theorem 2.1. Therefore the proof is finished once we show that $\gamma(B_X)$ is stronger than the weak-topology on W , or equivalently each member x^{**} of $B_{X^{**}}$ is continuous on $(W, \gamma(B_X))$. So fix an element x^{**} in $B_{X^{**}}$. By Corollary 4.4, $(H, \text{weak})^{\mathbb{N}}$ is Lindelöf, and therefore, by Theorem A, $(C(H, \text{weak}), \tau_p(H))$ is countably tight. Since $B_X|_H$ is $\tau_p(H)$ -dense in $B_{X^{**}}|_H \subset C(H, \text{weak})$, there is a countable subset $A \subset B_X$ such that $x^{**}|_H$ is in the $\tau_p(H)$ -closure of $A|_H$. Let G be the convex hull of H . Then by the linearity, $x^{**}|_W$ is in the $\tau_p(G)$ -closure of $A|_W$. By (a), G is norm-dense in W and $B_{X^{**}}|_W$ is an equicontinuous family of functions on $(W, \|\cdot\|)$. Hence $\tau_p(W)$ and $\tau_p(G)$ coincide on $B_{X^{**}}|_W$, and so $x^{**}|_W$ is in the $\tau_p(W)$ -closure of $A|_W$. Finally, $A|_W$ is an equicontinuous family on $(W, \gamma(B_X))$ and hence $x^{**}|_W$, being in the pointwise closure of $A|_W$, is $\gamma(B_X)$ -continuous on W . ■

Remark. In the theorem above as well as in the next corollary, the weak*-closed convex hull of H ($= \overline{\text{co}(H)}^{w^*}$) can be replaced by the weak*-closed absolutely convex hull of H . The proof is almost identical as above since [24, Theorem 2.5] is actually stated for the weak*-closed absolute convex hull case.

Corollary 4.6. *Let X be a Banach space, H a weak*-compact subset of X^* and W its weak*-closed convex hull. The following statements are equivalent*

- (a) (H, weak) is Lindelöf;
- (b) $(H, \text{weak})^{\mathbb{N}}$ is Lindelöf;
- (c) (W, weak) is Lindelöf;
- (d) $(W, \text{weak})^{\mathbb{N}}$ is Lindelöf.

Proof. - The implications (a) \Rightarrow (b) and (c) \Rightarrow (d) both follow from Corollary 4.4. The implications (b) \Rightarrow (a), (d) \Rightarrow (c) and (c) \Rightarrow (a) are obvious. And finally, the implication (a) \Rightarrow (c) is Theorem 4.5. ■

5. BANACH SPACES GENERATED BY RN-COMPACT SUBSETS

If X is either a weakly compactly generated Banach space or the dual of an Asplund space, then X is generated by an RN-compact subset in weak- or weak*-topology. We shall deal in this section with the class of Banach spaces generated by RN-compact subsets with respect to a topology weaker than the weak topology. Being more concrete, our framework is the following: for a Banach space $(X, \|\cdot\|)$ we consider $F \subset X^*$ a *norming subset* (also called *1-norming subset*) for X , that is, a \mathbb{Q} -linear set F satisfying

$$(5) \quad \|x\| = \sup\{|\langle x, f \rangle| : f \in F \cap B_{X^*}\}.$$

If a bounded set $H \subset X$ is $\sigma(X, F)$ -compact and fragmented by the norm, then $(H, \sigma(X, F))$ is an RN-compact set since the norm is $\sigma(X, F)$ -lower semicontinuous, and we will study the space generated by it, that is, the space $Y = \overline{\text{span}(H)}^{\|\cdot\|}$. The Banach space Y thus obtained will be called a *Banach space generated by an RN-compact subset*. In Section 7, we exhibit several examples of such Banach

spaces. In order to show the main properties of spaces generated this way we shall first see that these spaces admit projectional generators as defined below. Here our main reference shall be [11]. If A is a non-empty subset of a Banach space X , A^\perp denotes the subset $\{f \in X^* : f(x) = 0 \text{ for all } x \in A\}$ of X^* .

Definition 2. *Let X be a Banach space. A projectional generator on X is a countable-valued map $\varphi : F \rightarrow 2^X$ on a norming subset $F \subset X^*$ such that whenever $B \subset F$ is a \mathbb{Q} -linear set, we have*

$$(6) \quad \varphi(B)^\perp \cap \overline{B \cap B_{X^*}^{w*}} = \{0\}$$

According to the method developed in [27], [25] and [11], the existence of a projectional generator leads to the existence of a *projectional resolution of identity* (PRI for short) in the sense that follows. Given a Banach space X , the *density character* of X (denoted by $\text{dens}X$) is defined to be the least cardinality of a dense subset of X . Let μ be the least ordinal such that $|\mu| = \text{dens}X$, where $|\mu|$ denotes the cardinality of the ordinal μ . A PRI on X is a transfinite sequence $\{P_\alpha : \omega_0 \leq \alpha \leq \mu\}$ of linear projections in X satisfying the following conditions, where α and β are arbitrary ordinals in $[\omega_0, \mu]$. (a) $\|P_\alpha\| = 1$; (b) $\text{dens}P_\alpha(X) \leq |\alpha|$; (c) $P_\alpha P_\beta = P_\beta P_\alpha = P_{\min\{\alpha, \beta\}}$; (d) For each $x \in X$ and each limit ordinal α , $P_\beta(x) \rightarrow P_\alpha(x)$ in the norm as $\beta \uparrow \alpha$. Next proposition gathers the main properties of spaces with a projectional generator. In what follows ‘‘LUR norm’’ stands for ‘‘locally uniformly rotund (or convex) norm’’.

Each part of the following theorem is known, but they are not usually stated in the form we prefer in the present paper. We record it here for reference.

Theorem 5.1. *Let X be a Banach space with a projectional generator $\varphi : F \rightarrow 2^X$. Then the following statements hold.*

- (a) *X admits a PRI $\{P_\alpha : \omega_0 \leq \alpha \leq \mu\}$ such that $P_\alpha(X)$ has a projectional generator for each $\omega_0 \leq \alpha < \mu$;*
- (b) *X admits an equivalent LUR norm;*
- (c) *There is a linear continuous one-to-one operator $T : X \rightarrow c_0(\Gamma)$, for some set Γ ;*
- (d) *The Banach space X is $\gamma(X, F)$ -Lindelöf, where $\gamma(X, F)$ is the topology on X of uniform convergence on bounded countable subsets of F .*

Proof.- (a) With the projectional generator φ in X , a PRI $\{P_\alpha : \omega_0 \leq \alpha \leq \mu\}$ can be constructed, based on pairs of \mathbb{Q} -linear subsets (A_α, B_α) , $A_\alpha \subset X$ and $B_\alpha \subset F$ with $\varphi(B_\alpha) \subset A_\alpha$ and B_α norming for A_α , see proposition 6.1.7 and remark 6.1.8 of [11]; so, we have $\overline{B_\alpha \cap B_{X^*}^{w*}} \cap A_\alpha^\perp = \{0\}$ and P_α is the projection from X onto $\overline{A_\alpha}^{\|\cdot\|}$ with kernel B_α^\perp . The space $P_\alpha^*(X^*) = \overline{B_\alpha}^{w*}$ is identified with the dual of $P_\alpha(X) = \overline{A_\alpha}^{\|\cdot\|}$ and therefore $P_\alpha(X)$ also has a projectional generator defined on B_α by $\varphi_\alpha(f) = P_\alpha(\varphi(f))$, $f \in B_\alpha$. These observations complete the proof of (a).

(b) and (c) Here we use the induction argument encapsulated in [7, Theorem VII.1.8]. Let \mathcal{P} be the class of Banach spaces that admit a projectional generator. Then (a) shows that the hypothesis for \mathcal{P} in [7, Theorem VII.1.8] is satisfied. Therefore each member X of \mathcal{P} admits an equivalent LUR norm. If, in the proof of [7, Theorem VII.1.8], one uses [11, Proposition 6.2.2.] instead of Theorem VII.1.6 of [7], then one can also conclude that each member X of \mathcal{P} has property (c).

(d) The proof of Theorem A in [25] give us this result. ■

What remains of this section is devoted to proving that a Banach space generated by an RN-compact subset has a projectional generator and therefore enjoys the properties listed in Theorem 5.1

First we recall Simons' lemma below, [31].

Lemma 3. *Let $(z_n)_n$ be a uniformly bounded sequence in $\ell^\infty(C)$ and let W be its convex hull. If B is a subset of C such that for every sequence of positive numbers $(\lambda_n)_n$ with $\sum_{n=1}^\infty \lambda_n = 1$ there is $b \in B$ such that*

$$(7) \quad \sup\left\{\sum_{n=1}^\infty \lambda_n z_n(y) : y \in C\right\} = \sum_{n=1}^\infty \lambda_n z_n(b),$$

then

$$(8) \quad \sup_{b \in B} \{\limsup_{n \rightarrow \infty} z_n(b)\} \geq \inf_C \{\sup w : w \in W\}.$$

A subset of X^* is said to be *total* if its linear span is weak*-dense in X^* . Clearly a norming subset for X is a total subset of X^* .

Definition 3. *Let X be a normed space, $C \subset X$ a set and F a total subset in X^* . A subset $B \subset C$ is said to be an F -boundary for C if for every f in F there is a $b \in B$ such that $f(b) = \sup\{f(x) : x \in C\}$.*

In what follows, when F is a total norm closed subspace of X^* we consider the norm associated to F given by

$$p_F(x) = \sup\{|\langle x, f \rangle| : f \in F \cap B_{X^*}\},$$

for $x \in X$. Then the unit ball of $(X, p_F)^*$ is the set $\overline{F \cap B_{X^*}}^{w^*}$ and $(X, p_F)^*$ is the subspace $H = \bigcup_{n=1}^\infty n \overline{F \cap B_{X^*}}^{w^*}$ of X^* . Clearly $F \subset H$.

Proposition 5.2. *Let X be a normed space and let F be a total norm closed subspace of X^* . Let C be a bounded subset of X and $B \subset C$ an F -boundary for C such that (B, p_F) is separable. Then we have*

$$(9) \quad \overline{\text{co}(B)}^{p_F} = \overline{\text{co}(C)}^{\sigma(X, F)}.$$

Proof.- The proof is based on the ideas in [13](see also [12]). As we remarked, the dual of (X, p_F) is the subspace $H = \bigcup_{n=1}^\infty n \overline{G}^{w^*}$ of X^* , where $G = B_{X^*} \cap F$, and $F \subset H$. Hence we have

$$\overline{\text{co}(B)}^{p_F} \subset \overline{\text{co}(C)}^{p_F} = \overline{\text{co}(C)}^{\sigma(X, H)} \subset \overline{\text{co}(C)}^{\sigma(X, F)}.$$

Assume that the conclusion of the proposition is false. Then there exists an element $x_0 \in \overline{\text{co}(C)}^{\sigma(X, F)} \setminus \overline{\text{co}(B)}^{p_F}$. Then by the separation theorem, there is a functional $f \in H = (X, p_F)^*$ such that

$$f(x_0) > \alpha > \sup\{f(b) : b \in B\}.$$

By scaling we may assume that $f \in \overline{G}^{w^*}$. Let $U = \{g \in X^* : g(x_0) > \alpha\}$. Then U is convex weak*-open and $f \in \overline{G}^{w^*} \cap U \subset \overline{G \cap U}^{w^*}$. Now \overline{G}^{w^*} is equicontinuous on (X, p_F) and B contains a countable p_F -dense subset D . Therefore in \overline{G}^{w^*} the topology of pointwise convergence on B is identical with the topology of

pointwise convergence on D , and the latter is pseudo-metrizable. It follows that there is a sequence $\{z_n : n \in \mathbb{N}\}$ in $G \cap U$ such that $\lim_n z_n(b) = f(b)$ for each $b \in B$. Our assumption of F being norm-closed and B being an F -boundary of C implies that the sequence $(z_n)_n$ satisfies the hypothesis of Lemma 3. Hence by (8),

$$\alpha > \sup_{b \in B} f(b) \geq \inf_{c \in C} \{\sup w(c) : w \in \text{co}(\{z_n\})\}.$$

It follows that $\alpha > \sup_C w$ for some $w \in \text{co}(\{z_n\}) \subset G \cap U$. In particular, since $w \in U$, $w(x_0) > \alpha > \sup_C w$. On the other hand, since $x_0 \in \overline{\text{co}(C)}^{\sigma(X,F)}$ and, being in F , w is $\sigma(X, F)$ -continuous, $w(x_0) \leq \sup_C w$, contradicting the previous inequality. This proves the proposition. ■

The pointwise limit of a sequence of real-valued continuous functions is called a function of the first Baire class. More generally a function f from a topological space M into a normed space X is said to be *of the first Baire class* if there is a sequence of continuous functions $f_n : M \rightarrow X$ such that $(f_n)_n$ converges to f in (X^M, τ_p) . A multivalued map φ from the topological space M to the space of subsets of a topological space T is said to be *usco* if $\varphi(m)$ is a compact non-empty subset of T for each $m \in M$ and if φ is *upper-semicontinuous* in the sense that, whenever U is an open subset of T , $\{m \in M : \varphi(m) \subset U\}$ is open in M .

Ideas in [14], see also [29], allow us to modify Jayne-Rogers' selection theorem, [19], to our situation below.

Theorem 5.3. *Let M be a metric space, X a normed space and F a total norm closed subspace of X^* . Let H be a norm bounded $\sigma(X, F)$ -compact subset of X which is fragmented by the norm p_F . If ψ is an usco map from M to subsets of $(H, \sigma(X, F))$, then ψ has a first Baire selector f from M into (X, p_F) .*

Proof. - If we identify (X, p_F) with a subspace of $\ell^\infty(F \cap B_{X^*})$ and H with a weak* compact subset there, then we can apply the Remark 17 in [18] to obtain a selector f of ψ_H which is σ -discrete and of the first Borel class from F to $\ell^\infty(B \cap B_{X^*})$ (see Corollary 7 in [18]). Such a selector as a map from F into (X, p_F) is also σ -discrete of the first Borel class, and by Theorems 1 and 2 of [29] f is first Baire class from F into (X, p_F) (see also [14] and the remarks in the introduction of [29]). ■

We are now ready to prove below one of the main properties of the selectors obtained above: the result that follows is a counterpart to the one stated as Theorem 26 in [18], and it is in the setting of topologies of pointwise convergence on total sets.

Theorem 5.4. *Let X be a normed space and let F be a total norm closed subspace of X^* . Let H be a norm bounded $\sigma(X, F)$ -compact subset of X . Let $\psi_H : F \rightarrow 2^H$ be the multi-valued map given by*

$$\psi_H(f) = \{x \in H : f(x) = \sup_H f\}.$$

Then ψ_H has a selector of the first Baire class from $(F, \|\cdot\|)$ into (X, p_F) if, and only if, $(H, \sigma(X, F))$ is fragmented by p_F . Moreover, if $f : F \rightarrow H$ is such a selector of ψ_H , then we have

$$(10) \quad \overline{\text{co}(H)}^{\sigma(X,F)} = \overline{\text{co}(f(F))}^{p_F}.$$

Proof.- The arguments here are similar to the ones in [18, Theorem 26]. First it is easy to check that ψ_H is an usco-map from $(F, \|\cdot\|)$ into compact subsets of $(H, \sigma(X, F))$. If $(H, \sigma(X, F))$ is fragmented by p_F , then, by Theorem 5.3, ψ_H has a first Baire class selector $f : (F, \|\cdot\|) \rightarrow (X, p_F)$. Conversely assume that such a selector f exists. Let S be a $\|\cdot\|$ -closed and $\|\cdot\|$ -separable subspace of F , and consider the quotient normed space $(X/S^\perp, \|\cdot\|_S)$. Recall that the dual of $(X/S^\perp, \|\cdot\|_S)$ is isometric with \overline{S}^{w^*} and hence S is a norm closed total subspace of $(X/S^\perp, \|\cdot\|_S)^*$. Let $\pi_S : X \rightarrow X/S^\perp$ be the canonical quotient map and let p_S be the norm on X/S^\perp given by

$$(11) \quad p_S(\pi_S(x)) = \overline{p}_S(x) \stackrel{\text{def}}{=} \sup\{|g(x)| : g \in S \cap B_{X^*}\}$$

for each $x \in X$. Then $\pi_S(H)$ is a $\|\cdot\|_S$ -bounded, $\sigma(X/S^\perp, S)$ -compact subset of X/S^\perp , and $\pi_S(f(S))$ is an S -boundary for $\pi_S(H)$. Now let $f_k : F \rightarrow X$ be a sequence of $\|\cdot\| - p_F$ continuous maps such that for each $g \in F$, $f_k(g) \rightarrow f(g)$ in p_F . For each subset A of F , let

$$\Phi(A) = \bigcup_{k=1}^{\infty} f_k(A).$$

Then $f(\overline{A}^{\|\cdot\|}) \subset \overline{\Phi(A)}^{p_F}$ and $\Phi(A)$ is countable whenever A is. If D is a $\|\cdot\|$ -dense countable subset of S , then $f(S) = f(\overline{D}^{\|\cdot\|}) \subset \overline{\Phi(D)}^{p_F}$. Hence $f(S)$ is p_F -separable and so $\pi_S(f(S))$ is p_S -separable. It follows from Proposition 5.2 that

$$(12) \quad \overline{\text{co}(\pi_S(f(S)))}^{p_S} = \overline{\text{co}(\pi_S(H))}^{\sigma(X/S^\perp, S)}.$$

This shows in particular that, whenever S is a $\|\cdot\|$ -separable $\|\cdot\|$ -closed subspace of F , $\pi_S(H)$ is p_S -separable and hence H is \overline{p}_S -separable. Regarding H as a τ_p -compact subset of $[-m, m]^{F \cap B_{X^*}}$ with an appropriate $m > 0$, we see from Theorem 2.1 that $(H, \sigma(X, F))$ is fragmented by p_F .

Finally we show that (10) is a consequence of (12). For this it is sufficient to prove that for each $u \in X$, there is a $\|\cdot\|$ -separable $\|\cdot\|$ -closed subspace S of F such that

$$(13) \quad p_S\text{-dist}(\pi_S(u), \text{co}(\pi_S(f(S)))) \geq p_F\text{-dist}(u, \text{co}(f(S))).$$

For if $u \in \overline{\text{co}(H)}^{\sigma(X, F)}$ and if S is chosen as above, then since

$$\pi_S(u) \in \overline{\pi_S(\text{co}(H))}^{\sigma(X/S^\perp, S)}$$

we have, by (12), $0 = p_S\text{-dist}(\pi_S(u), \text{co}(\pi_S(f(S)))) \geq p_F\text{-dist}(u, \text{co}(f(S)))$. Hence $u \in \overline{\text{co}(f(S))}^{p_F} \subset \overline{\text{co}(f(F))}^{p_F}$. This shows that the left side of (10) is contained in the right side. The reverse inclusion is obvious.

To prove (13), let $u \in X$. For each countable subset M of X , let $\alpha(M)$ be a countable subset of $F \cap B_{X^*}$ such that, for each $x \in M$,

$$p_F(u - x) = \sup\{|g(u - x)| : g \in \alpha(M)\}.$$

Inductively we define a sequence $A_1 \subset A_2 \subset \dots$ of countable subsets of F as follows: let g_0 be an arbitrary non-zero element of F and let $A_1 = \{qg_0 : q \in \mathbb{Q}\}$.

Assuming that A_n has been defined, let

$$A_{n+1} = \text{span}_{\mathbb{Q}}(\alpha(\text{co}_{\mathbb{Q}}(\Phi(A_n))) \cup A_n),$$

where $\text{span}_{\mathbb{Q}}(C)$ (resp. $\text{co}_{\mathbb{Q}}(C)$) denotes the set of all linear (resp. convex) combinations of elements of C with rational coefficients. Let $S = \overline{\bigcup_{n=1}^{\infty} A_n}^{\|\cdot\|}$.

Before showing this S satisfies (13), we note that if $y \in \text{co}_{\mathbb{Q}}(\Phi(A_n))$ then $p_F(u - y) = \sup\{|g(u - y)| : g \in \alpha(\text{co}_{\mathbb{Q}}(\Phi(A_n)))\} \leq \bar{p}_S(u - y) \leq p_F(u - y)$. Hence $p_F(u - y) = \bar{p}_S(u - y)$. Now by the definition of Φ ,

$$\overline{\text{co}(f(S))}^{p_F} \subset \overline{\text{co}(\Phi(\bigcup_{n=1}^{\infty} A_n))}^{p_F} \subset \overline{\text{co}_{\mathbb{Q}}(\bigcup_{n=1}^{\infty} \Phi(A_n))}^{p_F} = \bigcup_{n=1}^{\infty} \overline{\text{co}_{\mathbb{Q}}(\Phi(A_n))}^{p_F}.$$

Let $x \in \text{co}f(S)$ and $\varepsilon > 0$ be arbitrary. Then there is a $y \in \text{co}_{\mathbb{Q}}(\Phi(A_n))$ for some n such that $\bar{p}_S(x - y) \leq p_F(x - y) < \varepsilon$. Then

$$\begin{aligned} p_S(\pi_S(u) - \pi_S(x)) &= \bar{p}_S(u - x) \geq \bar{p}_S(u - y) - \varepsilon = p_F(u - y) - \varepsilon \geq \\ &\geq p_F(u - x) - 2\varepsilon \geq p_F\text{-dist}(u, \text{co}f(S)) - 2\varepsilon. \end{aligned}$$

Since $x \in f(S)$ and $\varepsilon > 0$ are arbitrary, we obtain (13). ■

Remark In the setting of Theorem 5.4, (12) is now true whenever S is a $\|\cdot\|$ -closed subspace of F . This can be seen by applying Theorem 5.4 to the normed space X/S^{\perp} , the total subspace S of $(X/S^{\perp})^* = \bar{S}^{w*}$, the $\sigma(X/S^{\perp}, S)$ -compact set $\pi_S(H)$ and the selector $\pi_S \circ f|_S$ for the usco map $\psi_S : S \rightarrow 2^{\pi_S(H)}$. This remark is important in the proof of the next theorem.

Theorem 5.5. *Let X be a Banach space, F a norming subset of X^* and let H be a bounded $\sigma(X, F)$ -compact subset of X fragmented by the norm of X . Then the Banach space $Y = \overline{\text{span}(H)}^{\|\cdot\|}$ has a projectional generator.*

Proof.- We first prove the case $X = Y$. Since H is bounded, $\sigma(X, F)$ and $\sigma(X, \bar{F}^{\|\cdot\|})$ coincide on H . Hence we may assume that F is a $\|\cdot\|$ -closed norming subspace. Let $\psi_H : F \rightarrow 2^H$ be the set-valued map given by $\psi_H(g) = \{x \in H : g(x) = \sup_H g\}$ for each $g \in F$. Then by Theorem 5.4, ψ_H admits a selector $f : F \rightarrow H$ of the first Baire class from $(F, \|\cdot\|)$ into $(X, \|\cdot\|)$. Let $\{f_k\}$ be a sequence of continuous maps: $(F, \|\cdot\|) \rightarrow (X, \|\cdot\|)$ such that $f_k(g) \rightarrow f(g)$ in the norm for each $g \in F$, and we define the countable-valued map $\varphi : F \rightarrow 2^X$ by $\varphi(g) = \{f_k(g) : k \in \mathbb{N}\}$. We prove that φ is a projective generator (cf. Definition 2). So let B be a \mathbb{Q} -linear subset of F , and let $g \in \varphi(B)^{\perp} \cap \overline{B \cap B_{X^*}}^{w*}$. We must show that $g = 0$.

Let $S = \bar{B}^{\|\cdot\|} \subset F$, let $\pi_S : X \rightarrow X/S^{\perp}$ be the quotient map and let p_S be the norm defined on X/S^{\perp} by (11). Since $g \in \overline{S \cap B_{X^*}}^{w*}$, g defines a p_S -continuous linear functional \bar{g} on X/S^{\perp} by the formula: $\bar{g}(\pi_S(x)) = g(x)$ for each $x \in X$. Now by the definition of φ , $f(S) = f(\bar{B}^{\|\cdot\|}) \subset \overline{\varphi(B)}^{\|\cdot\|}$. Since g vanishes on $\varphi(B)$, it also vanishes on $f(S)$, and hence \bar{g} vanishes on $\pi_S(f(S))$. By the remark following the last theorem, (12) is valid for S and hence $\pi_S(H) \subset \overline{\text{co}(\pi_S(f(S)))}^{p_S}$. Therefore by continuity \bar{g} vanishes on $\pi_S(H)$, i.e. g vanishes on H . Since X is the norm-closed span of H , $g = 0$.

The general case is proved by applying the special case above to the Banach space Y and the norming subspace $F|_Y$ for Y . Note that H is a $\sigma(Y, F|_Y)$ -compact subset of Y and it is fragmented by the norm of Y . ■

Corollary 5.6. *Let X be a Banach space, F a norming subset of X^* , H a bounded subset of X which is $\sigma(X, F)$ -compact and fragmented by the norm of X , and let $Y = \overline{\text{span}(H)}^{\|\cdot\|}$. Then,*

- (a) $(Y, \gamma(X, F))$ is Lindelöf;
- (b) Y has a PRI;
- (c) Y has an equivalent LUR norm.

Proof.- This is a straightforward consequence of Theorem 5.1 and Theorem 5.5. ■

Another property of spaces generated by RN-compact sets is the following. For this, we need one more definition. Let (Z, τ) be a topological space and ρ a metric on Z . Then (Z, τ) is said to be σ -fragmented by ρ if for each $\varepsilon > 0$, Z can be written as $Z = \bigcup\{Z_n : n \in \mathbb{N}\}$ with each Z_n having the property that, whenever C is a non-empty subset of Z_n , there exists a τ -open subset U of Z such that $U \cap C$ is non-empty and of ρ -diameter less than ε .

Theorem 5.7. *Let X be a Banach space, F a norming subset of X^* , H a bounded subset of X which is $\sigma(X, F)$ -compact fragmented by the norm of X and let $Y = \overline{\text{span}(H)}^{\|\cdot\|}$. Then, $(Y, \sigma(X, F))$ is σ -fragmented by the norm.*

Proof.- The proof analogous to the one given for weakly compactly generated in [15]. Indeed, $W = \overline{\text{co}(H)}^{\sigma(X, F)} = \overline{\text{co}(H)}^{\|\cdot\|}$ is $\sigma(X, F)$ -compact and fragmented by the norm, [6, 4.1, 5.2 and 5.3]. Lemmas 2.1 and 2.2 [24] gives us that $W - W$ is again $\sigma(X, F)$ -compact and fragmented by the norm. We now have $Y = \overline{\bigcup_{n=1}^{\infty} n(W - W)}^{\|\cdot\|}$ and because F is norming, the norm in Y is $\sigma(X, F)$ -lower semi-continuous and Lemma 2.3 in [15] gives us the conclusion. ■

We can gather all the information that we have obtained so far in the following:

Theorem 5.8. *Let X be a Banach space, F a norming subset of X^* , H a bounded subset of X which is $\sigma(X, F)$ -compact and let $Y = \overline{\text{span}(H)}^{\|\cdot\|}$. The following statements are equivalent,*

- (a) $(H, \sigma(X, F))$ is fragmented by the norm;
- (b) $(Y, \sigma(X, F))$ is σ -fragmented by the norm;
- (c) $(H, \gamma(X, F))$ is Lindelöf;
- (d) $(Y, \gamma(X, F))$ is Lindelöf.

Proof.- (a) \Rightarrow (b) is the previous result. (b) \Rightarrow (a) follows from Lemma 3.1.1 in [16]. (a) \Rightarrow (d) is the item (a) of Corollary 5.6. (d) \Rightarrow (c) is obvious and (c) \Rightarrow (a) is also the implication (c) \Rightarrow (a) in Theorem 2.1. ■

In terms of compact sets embedded in cubes, the theorem above can be rephrased as:

Theorem 5.9. *Let $K \subset [-1, 1]^D \subset \ell^\infty(D)$ be a τ_p -compact set. The following statements are equivalent*

- (a) (K, τ_p) is fragmented by the norm;
- (b) $(\overline{\text{span}(K)}^{\|\cdot\|}, \tau_p)$ is σ -fragmented by the norm;

- (c) $(K, \gamma(D))$ is Lindelöf;
 (d) $(\overline{\text{span}(K)}^{\|\cdot\|}, \gamma(D))$ is Lindelöf.

6. BANACH SPACES GENERATED BY LINDELÖF SUBSETS

In this section we study Banach spaces which are Lindelöf in weak topology. Main tools are again the projectional generators. Beyond Theorem 6.1 below, that gives a quite general way of deciding when a Banach space is weakly Lindelöf, here we take advantage of the scope of the results in Section 4 and the main results in [4] to prove that a Banach space X generated by a weakly Lindelöf subset which is $\sigma(X, F)$ -compact with respect to some norming subspace $F \subset X^*$, is weakly Lindelöf. We need the following definition. For each set Γ , let $\Sigma(\Gamma)$ be the subspace of $\ell^\infty(\Gamma)$ consisting of all $u \in \ell^\infty(\Gamma)$ with $\{\gamma : u(\gamma) \neq 0\}$ at most countable. A compact Hausdorff space K is said to be *Corson* if, for some Γ , K can be embedded in $\Sigma(\Gamma)$ as a pointwise compact subset.

Definition 4 ([1]). *A Banach space X is said to be Weakly Lindelöf Determined (WLD for short) if there is a bounded one-to-one linear map $T : X^* \rightarrow \ell^\infty(\Gamma)$, for some set Γ , which is $\sigma(X^*, X)$ -pointwise continuous and such that $T(X^*) \subset \Sigma(\Gamma)$*

It was established in [26] that a Banach space is WLD if, and only if, its dual unit ball with the weak* topology is Corson compact. Note that WCG Banach spaces and hence separable Banach spaces are WLD. It is known that a WLD Banach space is $\gamma(B_{X^*})$ -Lindelöf, [25] and renormable by a LUR norm, [34] and [21]. A Banach space X or more generally a convex subset M of X is said to have *property C* (after Corson) if each collection of relatively closed convex subsets of M with empty intersection has a countable subcollection with empty intersection. If (M, weak) is Lindelöf, then M has property C since closed convex sets in X are also weak-closed. It is shown in [28] that the Banach space X has the property C if and only if, whenever $A \subset X^*$ and $f \in \overline{A}^{w^*}$, there is a countable subset C of A such that $f \in \overline{\text{co } C}^{w^*}$. This fact is crucial in the proof of the next theorem.

Theorem 6.1. *Let X be a Banach space with a projectional generator. If X has property C, then X is WLD, i.e. (B_{X^*}, weak^*) is Corson compact.*

Proof.- Let $\varphi : F \rightarrow 2^X$ be a projectional generator on X , where F is a norming subspace for X . Then X admits a PRI constructed as we have recalled in the Proposition 5.1. Let $\{P_\alpha : \omega_0 \leq \alpha \leq \mu\}$ be this PRI. Since property C is stable under taking closed subspaces, each $P_\alpha(X)$ has property C and a projectional generator. Now, by a standard induction process on the density character of the Banach space, we may assume that X admits a PRI $\{P_\alpha : \omega_0 \leq \alpha \leq \mu\}$, with μ a limit ordinal, such that, for each $\omega_0 \leq \alpha < \mu$, $P_\alpha(X)$ is WLD; that is, there is a one-to-one norm one operator

$$T_\alpha : P_\alpha^*(X^*) \rightarrow \ell^\infty(\Gamma_\alpha) \text{ with } T_\alpha(P_\alpha^*(X^*)) \subset \Sigma(\Gamma_\alpha)$$

which is weak*-pointwise continuous. Assume that $\{\Gamma_\alpha : \omega_0 \leq \alpha < \mu\}$ is a disjoint family. Then we define

$$\Gamma = \Gamma_{\omega_0} \cup \bigcup \{\Gamma_{\alpha+1} : \omega_0 \leq \alpha < \mu\}$$

and $T : X^* \rightarrow \ell^\infty(\Gamma)$ by the formulas

$$(Tf)(n) = T_{\omega_0}(P_{\omega_0}^*(f))(n) \text{ if } n \in \Gamma_{\omega_0} = \mathbb{N}$$

$$(Tf)(\gamma) = T_{\alpha+1}(P_{\alpha+1}^*(f) - P_{\alpha}^*(f))(\gamma) \text{ if } \gamma \in \Gamma_{\alpha+1}, \omega_0 \leq \alpha < \mu.$$

Clearly T is bounded linear and weak*-pointwise continuous. We claim that $T(X^*) \subset \Sigma(\Gamma)$. To prove it, we will see that the set $\{\alpha \in [\omega_0, \mu) : P_{\alpha+1}^*(f) - P_{\alpha}^*(f) \neq 0\}$ is at most countable for each $f \in X^*$. Assume on the contrary that this is not the case and take $f \in X^*$ so that this set is uncountable. Recall that the family $\{B_{\alpha} : \alpha < \mu\}$ is a long sequences of increasing \mathbb{Q} -linear subsets of F with $P_{\alpha}^*(X^*) = \overline{B_{\alpha}}^{w^*}$ for each $\alpha < \mu$. Also for each limit ordinal $\beta \leq \mu$ and $f \in X^*$, $\text{weak}^*\text{-}\lim_{\alpha \uparrow \beta} P_{\alpha}^*(f) = P_{\beta}^*(f)$, and $P_{\mu}^* = \text{Id}$. Let $\Delta = \{\alpha \in [\omega_0, \mu) : P_{\alpha+1}^*(f) - P_{\alpha}^*(f) \neq 0\}$. Then Δ is an uncountable subset of $[\omega_0, \mu)$ which is well-ordered under the inherited ordering. Therefore there is an order-isomorphism φ from $[0, \omega_1)$ onto an initial segment of Δ . Let $\eta = \sup \varphi([0, \omega_1)) \leq \mu$. Then $P_{\eta}^*(f) = \text{weak}^*\text{-}\lim_{\gamma \uparrow \omega_1} P_{\varphi(\gamma)}^*(f)$ and therefore

$$P_{\eta}^*(f) \in \overline{\bigcup_{\gamma < \omega_1} P_{\varphi(\gamma)}^*(f)}^{w^*}.$$

Since X has property \mathcal{C} , there is a sequence $\gamma_1 < \gamma_2 < \dots$ in $[0, \omega_1)$ such that

$$P_{\eta}^*(f) \in \overline{\text{co}(\bigcup_{i=1}^{\infty} P_{\varphi(\gamma_i)}^*(f))}^{w^*}.$$

Let $\xi = \varphi(\sup_i \gamma_i) \in \Delta$. Then $\xi < \eta \leq \mu$. Since for each i , $P_{\varphi(\gamma_i)}^*(f) \in \overline{B_{\varphi(\gamma_i)}}^{w^*} \subset \overline{B_{\xi}}^{w^*}$, we have $P_{\eta}^*(f) \in \overline{B_{\xi}}^{w^*}$. It follows that $P_{\eta}^*(f)$ is a fixed point of P_{α}^* for all $\alpha \geq \xi$. Hence if $\xi \leq \alpha < \eta$, then $P_{\eta}^*(f) = P_{\alpha}^*P_{\eta}^*(f) = P_{\alpha}^*(f)$ by the property of PRI: $P_{\eta}P_{\alpha} = P_{\min\{\eta, \alpha\}}$. In particular, $P_{\xi+1}^*(f) = P_{\eta}^*(f) = P_{\xi}^*(f)$, contradicting $\xi \in \Delta$. Hence $T(X^*) \subset \Sigma(\Gamma)$.

To see that T is one-to-one, let $T(f) = 0$ for an $f \in X^*$. Then $P_{\omega_0}^*(f) = 0$, and $P_{\alpha+1}^*(f) = P_{\alpha}^*(f) = 0$ for all $\alpha \in [\omega_0, \mu)$. Then by a straightforward (transfinite) induction, $P_{\alpha}^*(f) = 0$ for all $\alpha \in [\omega_0, \mu)$, and hence $f = \text{weak}^*\text{-}\lim_{\alpha \uparrow \mu} P_{\alpha}^*(f) = 0$. ■

A combination of Theorem 5.5 and Theorem 6.1 gives us the following:

Corollary 6.2. *Let X be a Banach space, F a norming subset of X^* , H a bounded subset of X which is $\sigma(X, F)$ -compact fragmented by the norm of X and let $Y = \overline{\text{span}(H)}^{\|\cdot\|}$. If Y has property \mathcal{C} , then Y is WLD.*

As mentioned earlier, a WLD Banach space is weakly Lindelöf, but its converse is not true; cf. [21, p. 514]. In [21, p. 521], Mercourakis and Negrepointis have asked if this converse is true in dual Banach spaces. The affirmative answer to this question is contained in [25] where it is shown that if X is an Asplund space then X^* is weakly Lindelöf if and only if $(B_{X^{**}}, \text{weak}^*)$ is Corson compact, i.e. X^* is WLD. Recall that Edgar had observed earlier [9] that X is an Asplund space whenever X^* is weakly Lindelöf. The next two corollaries are generalizations of the result in [25] just mentioned. The first one is a special case of the previous corollary.

Corollary 6.3. *Let X be an Asplund space, H a subset of X^* which is weak*-compact and let $Y = \overline{\text{span}(H)}^{\|\cdot\|}$. If Y has property \mathcal{C} then Y is WLD. In particular, if X is an Asplund space, then X^* is WLD if and only if it has property \mathcal{C} .*

A combination of most of the results in this paper and the main result in [4] finally allows us to prove.

Corollary 6.4. *Let X be a Banach space, H a subset of X^* which is weak*-compact and weakly Lindelöf. Then, the space generated by H , $Y = \overline{\text{span}(H)}^{\|\cdot\|}$, is WLD. In particular Y is weakly Lindelöf.*

Proof.- By the remark following Theorem 4.5, we know that the weak*-closed absolute convex hull of H , say W , is also weakly Lindelöf. Hence by Corollary E of [4], (W, weak) is fragmented by the norm. Furthermore, $Y = \overline{\text{span}(W)}^{\|\cdot\|} = \overline{\bigcup_{n=1}^{\infty} nW}^{\|\cdot\|}$ has property \mathcal{C} by Proposition 2 in [28]. Hence by Corollary 6.2, Y is WLD, and since a closed subspace of a WLD Banach space is again WLD ([21]), the corollary follows. ■

Remark 6.5. Let us remark that the statement Y being weakly Lindelöf in the previous Corollary can be proved more directly using Proposition 4.1, Corollary 5.6 and Theorem 4.5. With the notation above we know that $Z \stackrel{\text{def}}{=} \text{span}(H) = \bigcup_{n=1}^{\infty} nW$ is weakly Lindelöf, because it is a countable union of weakly Lindelöf subsets. Therefore by Proposition 4.1, $\gamma(X^*, X)$ is stronger than the weak topology on Z . On the other hand, $Y = \overline{Z}^{\|\cdot\|}$ is Lindelöf with respect to $\gamma(X^*, X)$ by Corollary 5.6, since H is weak*-compact and (H, weak) is fragmented by the norm. Consequently, Y will be weakly Lindelöf if we can prove that $\gamma(X^*, X)$ is stronger than the weak topology on Y . For this, it is sufficient to prove that for each $x^{**} \in X^{**}$ the restriction $x^{**}|_Y$ is $\gamma(X^*, X)$ -continuous. We know from above that $x^{**}|_Z$ is $\gamma(X^*, X)$ -continuous. This means that for each $\varepsilon > 0$ there is a $\gamma(X^*, X)$ -open neighborhood $U \subset X^*$ of the origin such that

$$(14) \quad |x^{**}(g)| < \varepsilon \quad \text{for each } g \in U \cap Z.$$

Now U is also $\|\cdot\|$ -open and therefore $\overline{U \cap Z}^{\|\cdot\|} = \overline{U \cap Y}^{\|\cdot\|} \supset U \cap Y$. Therefore the $\|\cdot\|$ -continuity of x^{**} and (14) imply that $|x^{**}(f)| \leq \varepsilon$ for every $f \in U \cap Y$. This means that x^{**} is $\gamma(X^*, X)$ -continuous on Y , which concludes the proof. ■

7. EXAMPLES OF SPACES GENERATED BY RN COMPACT SUBSETS

As mentioned at the beginning of Section 5, in this section we give several examples of Banach spaces generated by an RN-compact subset. By Theorem 5.5, these spaces possess all the properties stated in Theorem 5.1. Also by Corollary 6.2, for these spaces being WLD is equivalent to having property \mathcal{C} .

Example A. *Spaces with 1-norming Markusevich basis.*

Let us recall that a *Markusevich basis* or *M-Basis* of a Banach space X is a subset $\{(x_i, f_i) : i \in I\}$ of $X \times X^*$ such that

- (a) $\overline{\text{span}\{x_i : i \in I\}}^{\|\cdot\|} = X$;
- (b) $\bigcap_{i \in I} \text{Kernel}(f_i) = \{0\}$;

(c) $f_j(x_i) = \delta_{ij}, i, j \in I.$

Let us consider the subspace $F \stackrel{\text{def}}{=} \text{span}\{f_i\}$ which is a total subspace in X^* after condition (b). If $K \stackrel{\text{def}}{=} \{x_i : i \in I\} \cup \{0\}$ then it is easy to see that K is a $\sigma(X, F)$ -compact that is fragmented by the norm. Indeed, $\{x_i : i \in I\}$ is a $\sigma(X, F)$ -discrete set with 0 as its unique limit point. When F is norming, the M-basis is then called a *norming M-basis*. Therefore any Banach space with norming M-basis is generated by an RN-compact subset. The σ -fragmentability of spaces with a norming M-basis was first proved in the paper [17]; here, it is a consequence of Theorem 5.8.

Example B. *Spaces of continuous functions.*

Let K be a compact space and D a dense subset of K . If $H \subset C(K)$ is $\tau_p(D)$ -compact, uniformly bounded, fragmented by the supremum norm and separates the points of K , then $C(K)$ is generated by an RN-compact set. Indeed, in this case the norming subspace of $C(K)^*$ is $F = \text{span}\{\delta_x : x \in D\}$ and we observe that for every $n = 1, 2, \dots$ the set

$$H^n \stackrel{\text{def}}{=} \{f_1 \cdot f_2 \cdot \dots \cdot f_n : f_i \in H, i = 1, 2, \dots, n\}$$

is $\sigma(C(K), F)$ -compact and fragmented by the norm after Lemmas 2.1 and 2.2 in [24]. Now, $W = \bigcup_{n=1}^\infty (1/n)H^n \cup \{0\}$ is also $\sigma(C(K), F)$ -compact and σ -fragmented by the norm, hence fragmented, [16, Theorem 4.1]. On the other hand, the Stone-Weierstrass theorem gives us the equality $\overline{\text{span}(W)}^{\|\cdot\|} = C(K)$ and so $C(K)$ is generated by a $\sigma(C(K), F)$ -compact subset fragmented by the norm.

Example C. *Spaces of continuous functions defined on solid compact spaces and on compact spaces defined through adequate families of sets.*

Let I be a set and consider the cube $[0, 1]^I$ with the product topology. Given $x \in [0, 1]^I$ let us write

$$\text{supp}(x) \stackrel{\text{def}}{=} \{i \in I : x(i) \neq 0\}$$

and

$$\mathcal{F}(I) = \{x \in [0, 1]^I : \text{supp}(x) \text{ is finite}\}.$$

We claim that if $K \subset [0, 1]^I$ is a compact subset such that $K \cap \mathcal{F}(I)$ is dense in K (i.e. K is a special type of *Valdivia compact space*), then $C(K)$ is generated by an RN-compact subset. Indeed, let us write $D = K \cap \mathcal{F}(I)$ and $\pi_i : [0, 1]^I \rightarrow [0, 1]$ to denote the canonical projection onto the i -th coordinate, for each $i \in I$. Without loss of generality we can, and do, assume that for each $i \in I$ there is $x \in K$ such that $\pi_i(x) \neq 0$, because otherwise we can remove from the index set I the element i that is not needed to embed K in $[0, 1]^I$. Observe that $\{\pi_i : i \in I\}$ is $\tau_p(D)$ -discrete and that each $\tau_p(D)$ -neighborhood of 0 must contains all but at most finitely many $\{\pi_i : i \in I\}$; therefore $\{\pi_i : i \in I\} \cup \{0\}$ is $\tau_p(D)$ -compact, $\|\cdot\|_\infty$ -fragmented and separating the points of K . We use now Example B to conclude that $C(K)$ is generated by an RN-compact subset.

A compact space $K \subset [0, 1]^I$ is said to be *solid* if whenever $x \in K$ and $y \in [0, 1]^I$ are such that either $y_i = x_i$ or $y_i = 0$, for every $i \in I$, then $y \in K$. Obviously, if $K \subset [0, 1]^I$ is solid, then $K \cap \mathcal{F}(I)$ is dense in K and therefore $C(K)$ is generated by a RN-compact after our former reasoning.

A particular situation to which we can apply the above is when we deal with compact spaces defined through adequate families of sets. Following Talagrand, [33], if I is a non empty set, a family \mathcal{A} of subsets of I is called *adequate* if

- (a) If $A \in \mathcal{A}$ and $B \subset A$, then $B \in \mathcal{A}$;
- (b) $\{i\} \in \mathcal{A}$, for every $i \in I$;
- (c) If $A \subset I$ and every finite subset of A belongs to \mathcal{A} , then $A \in \mathcal{A}$.

If \mathcal{A} is an adequate family in I , then

$$K \stackrel{\text{def}}{=} \{\chi_A : A \in \mathcal{A}\}$$

is a solid compact space. Then $C(K)$ is also generated by a RN-compact subsets. Talagrand produced in [33, Théorème 4.3] an example of a compact space K defined through an adequate family of sets that is not Eberlein compact: the corresponding $C(K)$ then does not contain a $\tau_p(K)$ -compact subset separating the points of K , even though it contains a $\tau_p(D)$ -compact set (for certain dense $D \subset K$), $\|\cdot\|_\infty$ -fragmented subset separating the points of K .

Example D. *Spaces of Bochner integrable functions.*

Let $(X, \|\cdot\|)$ be a Banach space and $F \subset X^*$ a norming subspace. It was stated in [6, Corollary 4.3] that if $\sigma(X, F)$ -separable compact subsets of X are $\|\cdot\|$ -separable then the $\sigma(X, F)$ -compact (norm bounded) subsets H of X are $\|\cdot\|$ -fragmented; this is indeed a consequence of the equivalence between the first two statements in Theorem 2.1: write $D = F \cap B_{X^*}$ and consider $H \subset [-1, 1]^D$; given $A \subset D$ countable the set $H|_A \subset [-1, 1]^A$ is compact and metrizable, therefore separable; then there is a $\sigma(X, F)$ -compact and separable $S \subset H$ such that $S|_A = H|_A$; the restriction map $[-1, 1]^D \rightarrow [-1, 1]^A$ is continuous for the corresponding uniform metrics and therefore $H|_A$ is d_A -separable, because S is d_D -separable (S is $\|\cdot\|$ -separable).

The above observation is useful in finding more compact spaces “living” in Banach spaces and fragmented by the norm without being necessarily weakly compact.

Given a probability space (Ω, Σ, μ) we will denote by $L^p(\mu, X)$, $1 \leq p < +\infty$, the Banach space of μ -strongly measurable X -valued p -Bochner integrable functions $f : \Omega \rightarrow X$ normed by

$$\|f\|_p = \left(\int_{\Omega} \|f\|^p d\mu \right)^{\frac{1}{p}}$$

The dual $L^p(\mu, X)^*$ of $L^p(\mu, X)$ is a space of weak* measurable functions and the space $L^q(\mu, X^*)$, $1 = \frac{1}{p} + \frac{1}{q}$, which can be identified isometrically with a subspace of $L^p(\mu, X)^*$, is a norming subspace. So $\sigma' = \sigma(L^p(\mu, X), L^q(\mu, X^*))$ is a Hausdorff topology which is weaker than the weak topology of $L^p(\mu, X)$; these two topologies coincide if, and only if, X^* has the RNP [8, IV.1.1]. It was shown in [6, Example E] that every σ' -separable compact subset of $L^p(\mu, X)$ is norm separable. Therefore, every σ' -compact subset of $L^p(\mu, X)$ is fragmented by the norm. Thus we can apply the results in sections 5 and 6 to say for instance that if $H \subset L^p(\mu, X)$ is σ' -compact then the space $Y = \overline{\text{span}(H)}^{\|\cdot\|_p}$ has a PRI. This result is related to the main result of [3] which asserts the existence of a bounded one-to-one operator from $\overline{\text{span}(H)}^{\sigma'}$ into some $c_0(\Gamma)$ which is σ' -pointwise continuous.

REFERENCES

- [1] S. Argyros and S. Mercourakis, *On weakly Lindelöf Banach spaces*, Rocky Mountain J. Math. **23** (1993), no. 2, 395–446.
- [2] A.V. Arkhangel'skii, *Topological function spaces*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1992.
- [3] J. Batt and G. Schlüchtermann, *Eberlein compacts in $L_1(X)$* , Studia Math. **83** (1986), no. 3, 239–250.
- [4] B. Cascales, I. Namioka, and G. Vera, *The Lindelöf property and fragmentability*, Proc. Amer. Math. Soc. **128** (2000), no. 11, 3301–3309.
- [5] B. Cascales and J. Orihuela, *On pointwise and weak compactness in spaces of continuous functions*, Bull. Soc. Math. Belg. Sér. B **40** (1988), no. 3, 331–352.
- [6] B. Cascales and G. Vera, *Topologies weaker than the weak topology of a Banach space*, J. Math. Anal. Appl. **182** (1994), no. 1, 41–68.
- [7] R. Deville, G. Godefroy, and V. Zizler, *Smoothness and renormings in Banach spaces*, vol. 64, Pitman Monographs and Surveys in Pure and Applied Mathematics, 1993.
- [8] J. Diestel and J. Uhl, *Vector measures*, vol. 15, Math. Surveys. Amer. Math. Soc., 1977.
- [9] G. A. Edgar, *Measurability in a Banach space*, Indiana Univ. Math. J. **26** (1977), no. 4, 663–677.
- [10] R. Engelking, *General topology*, PWN- Polish Scientific Publishers., 1977.
- [11] M. Fabian, *Gâteaux differentiability of convex functions and topology*, John Wiley & Sons Inc., New York, 1997.
- [12] G. Godefroy, *Boundaries of a convex set and interpolation sets*, Math. Ann. **277** (1987), no. 2, 173–184.
- [13] ———, *Five lectures in geometry of Banach spaces*, Seminar on Functional Analysis, 1987 (Murcia, 1987), Univ. Murcia, Murcia, 1988, pp. 9–67.
- [14] R.W. Hansell, *On Borel mappings and Baire functions*, Trans. Amer. Math. Soc. **194** (1974), 195–211.
- [15] J. E. Jayne, I. Namioka, and C. A. Rogers, *σ -fragmentable Banach spaces*, Mathematika **39** (1992), no. 2, 197–215.
- [16] ———, *Topological properties of Banach spaces*, Proc. London Math. Soc. **66** (1993), 651–672.
- [17] ———, *σ -fragmented Banach spaces. II*, Studia Math. **111** (1994), no. 1, 69–80.
- [18] J. E. Jayne, J. Orihuela, A. J. Palleares, and G. Vera, *σ -fragmentability of multivalued maps and selection theorems*, J. Funct. Anal. **117** (1993), no. 2, 243–273.
- [19] J. E. Jayne and C. A. Rogers, *Borel selectors for upper semicontinuous set-valued maps*, Acta Math. **155** (1985), no. 1-2, 41–79.
- [20] J. L. Kelley, *General topology*, Springer-Verlag, New York, 1975.
- [21] S. Mercourakis and S. Negrepointis, *Banach spaces and topology. II*, Recent progress in general topology (Prague, 1991), North-Holland, Amsterdam, 1992, pp. 493–536.
- [22] P. R. Meyer, *The Baire order problem for compact spaces*, Duke Math. J. **33** (1966), 33–39.
- [23] ———, *Function spaces and the Aleksandrov-Uryshon conjecture*, Ann. Mat. Pura and Appl. **86** (1970), 25–29.
- [24] I. Namioka, *Radon-Nikodym compact spaces and fragmentability*, Mathematika **34** (1989), no. 2, 258–281.
- [25] J. Orihuela, *On weakly Lindelöf Banach spaces*, Progress in Functional Analysis Math. Studies North Holland **170** (1992), 279–291.
- [26] J. Orihuela, W. Schachermayer, and M. Valdivia, *Every Radon-Nikodym Corson compact space is Eberlein compact*, Studia Math. **98** (1991), no. 2, 157–174.
- [27] J. Orihuela and M. Valdivia, *Projective generators and resolutions of identity in Banach spaces*, Rev. Mat. Univ. Complutense Madrid **2** (1989), 179–199.
- [28] R. Pol, *On a question of H.H. Corson and some related problems*, Fund. Math. **109** (1980), no. 2, 143–154.
- [29] C. A. Rogers, *Functions of the first Baire class*, J. London Math. Soc. (2) **37** (1988), no. 3, 535–544.
- [30] Z. Semadeni, *Banach spaces of continuous functions. Vol. I*, PWN—Polish Scientific Publishers, Warsaw, 1971, Monografie Matematyczne, Tom 55.

- [31] S. Simons, *A convergence theorem with boundary*, Pacific J. Math. **40** (1972), 703–708.
- [32] M. Talagrand, *Deux generalisations d'un théorème de I.Namioka*, Pacific J. Math. **81** (1979), no. 1, 239–251.
- [33] ———, *Espaces de Banach faiblement K -analytiques*, Ann. of Math. **110** (1979), 407–438.
- [34] M. Valdivia, *Quasi-LB-spaces*, J. London Math. Soc. (2) **35** (1987), no. 1, 149–168.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, 30.100 ESPINARDO
MURCIA, SPAIN

E-mail address: beca@um.es

DEPARTMENT OF MATHEMATICS, BOX 354350, UNIVERSITY OF WASHINGTON, SEATTLE,
WASHINGTON 98195–4350, U.S.A.

E-mail address: namioka@math.washington.edu

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE MURCIA, 30.100 ESPINARDO
MURCIA, SPAIN

E-mail address: joseori@um.es