

# Spatio-temporal dynamic modeling of plant communities responses to hydrological pressures in a semiarid Mediterranean wetland

J. Martínez-López<sup>1</sup>, J. Martínez-Fernández<sup>1,2</sup>, B. Naimi<sup>3</sup>, M.F. Carreño<sup>1</sup> and M.A. Esteve<sup>1</sup>

<sup>1</sup>Ecology and Hydrology Department - University of Murcia (Murcia, Spain)

<sup>2</sup>Applied Biology Dept. University Miguel Hernandez (Elche, Spain)


<sup>3</sup>ITC - University of Twente (Enschede, The Netherlands)

October 28th - 31st



isem 2013  
TOULOUSE - FRANCE

# Study area



# Marina del Carmoli wetland (300 ha)

Wetland model

ISEM 2013

Introduction

Methods

Results

Conclusions



# Wetland plant communities

Semiarid Mediterranean saline wetlands are semi-terrestrial ecosystems

## HUMIDITY




## SALINITY

- ▶ Salt steppe (left) - priority habitat by the Habitats Directive
- ▶ Salt marsh (center) - habitat of interest by the HD
- ▶ Reed beds (right) (*Phragmites australis*) - invasive


# External water inputs

Percentage of irrigated areas has increased in the last decades due to the opening of a water transfer (Martínez-López et al., in press)



# Plant communities change

Important plant communities are being lost!



Carreño et al., 2008; Martínez-López et al., 2012

Introduction

Methods

Results

Conclusions

# Objective

- ▶ Spatially explicit wetland model of how irrigated agriculture is affecting plant community composition in this semiarid Mediterranean wetland

# Modelling environment


- ▶ R as a modelling environment:
  - ▶ GIS capabilities
  - ▶ source code is flexible
  - ▶ free availability and growing user community






# State variables

- ▶ Wetland is divided into pixels (25 m)
- ▶ Plant communities are modelled separately pixel by pixel (4 maps)
- ▶ The total abundance of plant communities within a pixel is limited so:
  - ▶ competition among plant communities mediated by
    - ▶ total drainage water input to the wetland
    - ▶ spatial environmental variables influencing water availability and growth
  - ▶ the dispersion of other PC from the surrounding pixels



# Initial and validation maps of plant communities

Model was tested by means of remote sensing data for the period 1992-2008




Carreño et al., 2008; Martínez-López et al., 2012


# Model assumptions I

- ▶ Increasing water input
- ▶ Only conversion to more humid / less saline plant communities

HUMIDITY




SALINITY



# Model assumptions II

## native vs. invasive taxa


- ▶ invasive reed beds are potentially present in all pixels
- ▶ salt marsh is able to disperse into neighbour pixels



# Non spatial forcing input


## Drainage water input

WARP index (Martínez-López et al., 2014a,b)




# Wetland environmental spatial parameters

- ▶ (A) distance map to ephemeral river 1 (**reed beds**)
- ▶ (B) distance map to ephemeral river 2 (**reed beds**)
- ▶ (C) Flow accumulation map (**salt marsh**)



- ▶ All parameters are on a relative 0–1 scale.

# Model diagram




# Model development/execution

1. Initial dynamic model was developed using Stella (1 pixel)
2. Conversion to R using 'StellaR' script (Naimi and Voinov, 2012)
3. State variables and spatial environmental variables as matrices
4. Model wrapped as a R function
5. `ode.2D("euler" method, time = 24 year, TS = 0.25)` (library "deSolve")



Overall Accuracy: 54% – 71%



1. The model serves as a tool for
  - ▶ wetland conservation and management studies (habitat loss)
  - ▶ testing plant community interactions
  - ▶ testing relationships between plant communities and environmental variables in space and time
2. Needs further developments
3. Source code will be documented and available on GitHub

# Thank you!